

```

CROSSTABS
  /TABLES=VAR06 BY VAR05 VAR07 VAR08
  /FORMAT= AVALUE TABLES
  /STATISTIC=CHISQ CORR
  /CELLS= COUNT COLUMN
  /COUNT ROUND CELL .

```

Crosstabs

Notes

Output Created		27-MAR-2009 07:27:26
Comments		
Input	Data	C:\My Documents\Benoit\Sites Web\EUT3018\spss\restaurants_categories.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	12
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		<pre> CROSSTABS /TABLES=VAR06 BY VAR05 VAR07 VAR08 /FORMAT= AVALUE TABLES /STATISTIC=CHISQ CORR /CELLS= COUNT COLUMN /COUNT ROUND CELL . </pre>
Resources	Elapsed Time	0:00:00,17
	Dimensions Requested	2
	Cells Available	174876
	Processor Time	0:00:00,02

[DataSet1] C:\My Documents\Benoit\Sites Web\EUT3018\spss\restaurants_categories.sav

Case Processing Summary

	Cases
--	-------

	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Ventes (Catégories) * Diplôme ITHQ	12	100,0%	0	,0%	12	100,0%
Ventes (Catégories) * Publicité (Catégories)	12	100,0%	0	,0%	12	100,0%
Ventes (Catégories) * Expérience (Catégories)	12	100,0%	0	,0%	12	100,0%

Ventes (Catégories) * Diplôme ITHQ

Crosstab

			Diplôme ITHQ		Total
			Non	OUI	Non
Ventes (Catégories)	400 K\$ ou moins	Count	3	0	3
		% within Diplôme ITHQ	100,0%	,0%	25,0%
	401 K\$ à 600 K\$	Count	0	5	5
		% within Diplôme ITHQ	,0%	55,6%	41,7%
	601 K\$ à 800 K\$	Count	0	3	3
		% within Diplôme ITHQ	,0%	33,3%	25,0%
	801 K\$ ou plus	Count	0	1	1
		% within Diplôme ITHQ	,0%	11,1%	8,3%
Total		Count	3	9	12
		% within Diplôme ITHQ	100,0%	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12,000(a)	3	,007
Likelihood Ratio	13,496	3	,004
Linear-by-Linear Association	6,195	1	,013
N of Valid Cases	12		

a 8 cells (100,0%) have expected count less than 5. The minimum expected count is ,25.

Symmetric Measures

		Value	Asymp. Std. Error(a)	Approx. T(b)	Approx. Sig.
Interval by Interval	Pearson's R	,750	,095	3,591	,005(c)
Ordinal by Ordinal	Spearman Correlation	,792	,125	4,108	,002(c)
N of Valid Cases		12			

a Not assuming the null hypothesis.

b Using the asymptotic standard error assuming the null hypothesis.

c Based on normal approximation.

Ventes (Catégories) * Publicité (Catégories)

Crosstab

			Publicité (Catégories)					Total 1000 \$ ou moins
			1000 \$ ou moins	1001 \$ à 2000 \$	2001 \$ à 3000 \$	3001 \$ à 4000 \$	4001 \$ ou plus	
Ventes (Catégories)	400 K\$ ou moins	Count	3	0	0	0	0	3
		% within Publicité (Catégories)	100,0%	,0%	,0%	,0%	,0%	25,0%
	401 K\$ à 600 K\$	Count	0	4	1	0	0	5
		% within Publicité (Catégories)	,0%	100,0%	100,0%	,0%	,0%	41,7%
	601 K\$ à 800 K\$	Count	0	0	0	3	0	3
		% within Publicité (Catégories)	,0%	,0%	,0%	100,0%	,0%	25,0%
	801 K\$ ou plus	Count	0	0	0	0	1	1
		% within Publicité (Catégories)	,0%	,0%	,0%	,0%	100,0%	8,3%
Total		Count	3	4	1	3	1	12
		% within Publicité (Catégories)	100,0%	100,0%	100,0%	100,0%	100,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	36,000(a)	12	,000
Likelihood Ratio	30,360	12	,002
Linear-by-Linear Association	10,411	1	,001
N of Valid Cases	12		

a 20 cells (100,0%) have expected count less than 5. The minimum expected count is ,08.

Symmetric Measures

		Value	Asymp. Std. Error(a)	Approx. T(b)	Approx. Sig.
Interval by Interval	Pearson's R	,973	,014	13,291	,000(c)
Ordinal by Ordinal	Spearman Correlation	,981	,023	16,062	,000(c)
N of Valid Cases		12			

a Not assuming the null hypothesis.

b Using the asymptotic standard error assuming the null hypothesis.

c Based on normal approximation.

Ventes (Catégories) * Expérience (Catégories)

Crosstab

			Expérience (Catégories)				Total
			1 à 3 ans	4 ou 5 ans	6 ou 7 ans	8 ou 9 ans	1 à 3 ans
Ventes (Catégories)	400 K\$ ou moins	Count	3	0	0	0	3
		% within Expérience (Catégories)	100,0%	,0%	,0%	,0%	25,0%
	401 K\$ à 600 K\$	Count	0	5	0	0	5
		% within Expérience (Catégories)	,0%	100,0%	,0%	,0%	41,7%
	601 K\$ à 800 K\$	Count	0	0	2	1	3
		% within Expérience (Catégories)	,0%	,0%	100,0%	50,0%	25,0%
	801 K\$ ou plus	Count	0	0	0	1	1
		% within Expérience (Catégories)	,0%	,0%	,0%	50,0%	8,3%
Total	Count	3	5	2	2	12	
	% within Expérience (Catégories)	100,0%	100,0%	100,0%	100,0%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	28,000(a)	9	,001
Likelihood Ratio	27,587	9	,001
Linear-by-Linear Association	10,241	1	,001
N of Valid Cases	12		

a 16 cells (100,0%) have expected count less than 5. The minimum expected count is ,17.

Symmetric Measures

		Value	Asymp. Std. Error(a)	Approx. T(b)	Approx. Sig.
Interval by Interval	Pearson's R	,965	,024	11,619	,000(c)
Ordinal by Ordinal	Spearman Correlation	,988	,015	20,602	,000(c)
N of Valid Cases		12			

a Not assuming the null hypothesis.

b Using the asymptotic standard error assuming the null hypothesis.

c Based on normal approximation.